

Assessment Of Nutritional Status In Hyperthyroidism Patients

Dr. Purnima Dey Sarkar*, Lincy K. Skaria**, Gopinath Agnihotram**

*Professor and HoD, **Research Scholar, Dept. of Biochemistry, MGM Medical College, Indore, M.P

Abstract : Background: Hyperthyroidism often referred to as overactive thyroid, is a type of thyrotoxicosis, a hypermetabolic clinical syndrome which occurs when there are elevated serum levels of T3 and/or T4. There are very less studies on the nutritional deficiencies related to hyperthyroidism. Methods: Quantitative determination of Thyrotropin (TSH), triiodothyronine (T3) and tetraiodothyronine (T4) was analyzed along with nutritional analysis of dietary nutrients like carbohydrate, proteins, fat, vitaminA, B1, B2, C, niacin, etc. Biostatistical analysis and correlations were analyzed by using Graph Pad prism software. Results: The mean intake of nutrients in Hyperthyroid and control groups are: Carbohydrate (147.9 ± 41.89 g, 218.3 ± 100.0 g), Thiamin (0.567 ± 2.850 mg, 1.051 ± 0.470 mg), Riboflavin (0.590 ± 0.341 mg, 1.121 ± 0.415 mg), Niacin (7.560 ± 2.346 mg, 11.59 ± 3.571 mg), pyridoxine (0.066 ± 0.262 mg, 0.317 ± 0.627 mg), Vitamin B1 (0.087 ± 0.205 µg, 0.199 ± 0.306 µg), Folicin (130.0 ± 36.57 µg, 143.4 ± 57.29 µg) and the mean values of all these nutrients were not optimum as recommended by ICMR. Conclusion: We should all be aware of the nutritional deficiencies encountered in hyperthyroidism and hence we can provide considerable support by eliminating adverse influences and normalizing the nutritional status by including multivitamins, antioxidants, etc. along with daily antithyroid drugs. [Skaria L et al NJIRM 2013; 4(4) : 8-11]

Key Words: Hyperthyroidism, Thyroid Stimulating Hormone (TSH), triiodothyronine (T3), tetraiodothyronine (T4), nutritional status, multivitamins.

Author for correspondence: Lincy K. Skaria, Siddhpur Dental College, Dethali, Siddhpur, Patan dist.

E-mail Address: linsey@rediffmail.com

Introduction: Hyperthyroidism, often referred to as an overactive thyroid, is a condition in which the thyroid gland produces and secretes excessive amounts of the free (not protein bound, and circulating in the blood¹) thyroid hormones, triiodothyronine (T3) and/or thyroxine (T4). Hyperthyroidism is a type of thyrotoxicosis, a hypermetabolic clinical syndrome which occurs when there are elevated serum levels of T3 and/or T4². Hyperthyroidism may be asymptomatic, but when it is not, symptoms are due to an excess of thyroid hormone. Therefore, some of the symptoms of hyperthyroidism are nervousness, irritability, increased perspiration, heart racing, hand tremors, anxiety, difficulty sleeping, thinning of the skin, fine brittle hair, and muscular weakness, weight loss, menstrual disturbances in women, etc.

Worldwide, approximately 1.1% population suffers from Hyperthyroidism³. An estimated 108 million people suffer from endocrine and Metabolic disorders in India and out of these 42 million suffer from thyroid disorders⁴. In Bastar region of Chattisgarh in India a study conducted on 150 subjects for thyroid disorders based on signs and symptoms confirmed that 59.3% were hypothyroid, 20.6% were euthyroid, 12.6% had

subclinical hypothyroidism, 7.3% were hyperthyroid⁵. Very less work on studies related to the assessment of nutritional status in Hyperthyroidism has been carried out, hence the study was conducted to evaluate the nutritional status in patients of hyperthyroidism.

Materials and Method: The subjects who visited the medicine department of M. G. M. medical college, Indore were carefully examined. 250 subjects were included in the study group of which 100 patients were of Hyperthyroid along with 150 controls. The quantitative determination of thyrotrophin (TSH), total triiodothyronine (T3) and total tetraiodothyronine (T4) in human serum was estimated by a microplate immunoenzymetric assay using the reagent kit by Monobind, Lake forest, USA^{6,7,8,9,10}. Each patient was in detail nutritionally analyzed by taking down the questionnaire along with 24 hour dietary recall proforma prepared and were nutritionally analyzed along with other nutrients like proteins, carbohydrates, fat, carotene, vitamin A, B1, folicin, thiamin, riboflavin, pyridoxine, calcium, phosphorous, sodium, magnesium, potassium, zinc¹¹.

Statistical analysis: Biostatistical analysis as well as correlations among parameters in individual groups were established by using Graph Pad Prism Software.

Results: There were two groups in the study group. The first group consisted of Hyperthyroid of 100 patients and second group consisted of control of 150 subjects.

Table 1 shows the distribution of Hyperthyroid patients in different levels of age. Hyperthyroidism was found to be increased among patients 20-to-49 years of age and declined with more than 50 years of age.

Table 1: Distribution table of Hyperthyroid patients in different levels of age groups.

Age groups	Hyperthyroidism (n=100)	Control (n=150)
10-19	07	14
20-29	26	31
30-39	25	54
40-49	28	27
50-59	12	17
60-69	02	07

Table 2 shows the difference of TSH, T3 and T4 levels in Hyperthyroid group compared to the control group. The mean value of TSH in Hyperthyroid group was $0.194 \pm 0.047 \mu\text{IU/mL}$ and in control was $5.082 \pm 1.571 \mu\text{IU/mL}$ and the difference was found to be significant ($p < 0.001$). The mean value of T4 in Hyperthyroid group was $14.63 \pm 2.631 \mu\text{g/dL}$ and in control was $8.333 \pm 1.827 \mu\text{g/dL}$ and the difference was found to be significant ($p < 0.001$).

Table 2: Mean and Standard deviation in Hyperthyroid group compared to the controls at T3, T4 and TSH levels.

Hormone	Hyperthyroidism (n=100)	Control (n=150)	p value
TSH	$0.194 \pm 0.047 \mu\text{IU/mL}$	$5.082 \pm 1.571 \mu\text{IU/mL}$	<0.001
T4	$14.63 \pm 2.631 \mu\text{g/dL}$	$8.333 \pm 1.827 \mu\text{g/dL}$	<0.001
T3	$2.608 \pm 0.891 \text{ ng/mL}$	$1.063 \pm 0.581 \text{ ng/mL}$	<0.001

The mean value of T3 levels in Hyperthyroid group was $2.608 \pm 0.891 \text{ ng/mL}$ and in control group was $1.063 \pm 0.581 \text{ ng/mL}$ and the difference was found to be significant ($p < 0.001$).

Table 3 shows correlation table of TSH, T3 and T4 levels in Hyperthyroid and control groups. There was a significant positive correlation between T3 and T4 in control group ($r = 0.5076$). Graph 1 shows the correlation between T3 and T4 levels in the control group.

Table 3: Correlation table of TSH, T4 and T3 levels in Hyperthyroid and control groups.

Correlation between hormones	Hyperthyroidism (n=100) r value	Control (n=150) r value
T3 & T4	0.1517 *	0.5076 **
T3 & TSH	-0.0380*	-0.0078 *
T4 & TSH	0.04007*	0.0629 *

* = Insignificant, ** = Significant positive correlation, ***= Significant negative correlation

Graph 1: Correlation table of T4 and T3 levels
Correlation between T₄ and T₃ in control




Table 4 shows the Mean values of Nutrient Intake by the subjects in Hyperthyroid and control group. The mean energy intake in Hyperthyroid and control groups were ($1517 \pm 337.8 \text{ kcal}$) and ($2198 \pm 393.6 \text{ kcal}$) respectively.

The mean intake of nutrients in Hyperthyroid and control groups are: Carbohydrate ($147.9 \pm 41.89 \text{ g}$, $218.3 \pm 100.0 \text{ g}$), Thiamin ($0.567 \pm 2.850 \text{ mg}$, $1.051 \pm 0.470 \text{ mg}$), Riboflavin ($0.590 \pm 0.341 \text{ mg}$, $1.121 \pm 0.415 \text{ mg}$), Niacin ($7.560 \pm 2.346 \text{ mg}$, $11.59 \pm 3.571 \text{ mg}$), pyridoxine ($0.066 \pm 0.262 \text{ mg}$,

0.317 ± 0.627 mg), Vitamin B1 (0.087 ± 0.205 µg, 0.199 ± 0.306 µg), Folic acid (130.0 ± 36.57 µg, 143.4 ± 57.29 µg) and the mean values of all these nutrients were not optimum as recommended by ICMR.

The mean intake of Zinc in Hyperthyroid group was (1.180 ± 2.904 mg) was very low compared to 10mg/day as recommended by ICMR. The mean values of Vitamin C in Hyperthyroid group was (102.8 ± 173.6 mg) was optimum as suggested by ICMR. While the mean values Sodium in hyperthyroid (94.40 ± 163.2 mg), Potassium (148.3 ± 252.1 mg), Magnesium (20.74 ± 25.79 mg) were not near the optimum as suggested by ICMR.

Table 4: Mean Nutrient Intake of subjects in Hyperthyroid and control groups.

Nutrient intake	Hyperthyroidism (n=100)	Control (n=150)
Protein	31.16 ± 12.44 g	57.07 ± 22.07 g
Fat	71.09 ± 22.85 g	101.3 ± 50.97 g
Saturated fat	25.29 ± 10.90 g	38.61 ± 9.935g
PUFA	9.330 ± 7.080 g	36.6 ± 16.54 g
Fibre	3.250 ± 9.230 g	4.707 ± 6.013 g
Carbohydrate	147.9 ± 41.89 g	218.3 ± 100.0 g
Energy	1517 ± 337.8 kcal	2198 ± 393.6 kcal
Calcium	313.4 ± 114.5 mg	411.2 ± 142.3 mg
Phosphorous	589.3 ± 227.9 mg	581.3 ± 176.7 mg
Carotene	241.4 ± 137.1 µg	460.3 ± 148.3 µg
Thiamin	0.567 ± 2.850 mg	1.051 ± 0.470 mg
Riboflavin	0.590 ± 0.341mg	1.121 ± 0.415 mg
Niacin	7.560 ± 2.346 mg	11.59 ± 3.571mg
Pyridoxine	0.066 ± 0.262 mg	0.317 ± 0.627 mg
Vitamin B1	0.087 ± 0.205 µg	0.199 ± 0.306 µg
Folic acid	130.0 ± 36.57 µg	143.4 ± 57.29 µg
Vitamin C	102.8 ± 173.6 mg	153.6 ± 54.50 mg
Zinc	1.180 ± 2.904 mg	9.240 ± 6.652 mg
Vitamin A	350.8 ± 1033 RE	403.7 ± 476.3 RE
Sodium	94.40 ± 163.2 mg	155.7 ± 238.2 mg
Potassium	148.3 ± 252.1 mg	456.9 ± 208.3 mg
Magnesium	20.74 ± 25.79 mg	28.07 ± 12.67 mg

Discussion: The calorie intake and the carbohydrate intake in hyperthyroid group were low compared with the controls which was similar to the study¹² where the dietary calorie and the carbohydrate intake in groups of hypothyroid and hypothyroid associated infertility was found to be low when compared with that of controls.

Also the mean value of nutrients like thiamin, Riboflavin, Niacin, Pyridoxine, Vitamin B1, Folic acid were found to be low in hyperthyroid group when compared with the control group which was similar to the study conducted in Hypothyroid patients the mean value of nutrients like thiamin, Riboflavin, Niacin, Pyridoxine, Vitamin B1, Folic acid were not optimum as suggested by the ICMR¹².

Also the zinc level was low in hyperthyroid group as suggested by the ICMR recommendation which again is similar to the study where zinc levels were low in hypothyroid groups¹². The possible explanation would be because of high phytate content due to consumption of goitrogenic foods seen among the patients of both the groups. Zinc effects on thyroid hormones are complex and include both synthesis and mode of action¹³. Thyroid transcription factors which are essential for modulation of gene expression contain zinc at cysteine residues¹⁴.

Thyroid conditions, especially hyperthyroidism, are characterized by serious nutritional deficiencies. Nutritional deficiencies cause a disruption of both endocrine and immune functions. The endocrine and immune systems are so closely related because their function rely upon the same essential nutrients.

The foods rich in high in B-vitamins and iron, such as whole grains, fresh cruciferous vegetables and sea vegetables, antioxidant foods including fruits such as blue berries, cherries, and tomatoes, must be included in hyperthyroid subjects. Refined foods, such as white breads, pastas, and sugar should be avoided.

Conclusion: The mean value of dietary intake of most of the nutrients in Hyperthyroid subjects were deficient and were not in accordance as suggested by the ICMR. We should all be aware of the nutritional deficiencies encountered in hyperthyroidism and hence we can provide considerable support by eliminating adverse influences and normalizing the nutritional status. Along with antithyroid drugs for the treatment of hyperthyroid subjects a multivitamin daily, containing the antioxidant vitamins A, C, E, the B-

complex vitamins, and trace minerals such as magnesium, calcium, zinc, and selenium should be included. Omega-3 fatty acids, such as fish oil, 1 - 2 capsules to help decrease inflammation and help with immunity. Vitamin C, 500 - 1,000 mg daily, as an antioxidant and for immune support. Alpha-lipoic acid, 25 - 50 mg twice daily, for antioxidant support.

References

1. Dario M. Torre, Geoffrey C. Lamb, Jerome Van Ruiswyk: Kochar's Clinical Medicine for Students. pp.402-403: Lippincott Williams & Wilkins; Fifth edition (2008) ISBN 0-7817-6699-0.
2. Kittisupamongkol W. Hyperthyroidism or thyrotoxicosis? Cleve Clin J Med. Mar 2009;76(3):152.
3. Sachdev Y. Hyperthyroidism A Subtle Disorder. Adapted from the Pharmaceutical Journal 2007; 265 (7109): 240-244.
4. Kochupillai N. Clinical Endocrinology in India. Curr Sci 2000 ; 79:1061-7.
5. Lincy K. Skaria, Purnima Dey Sarkar, Gopinath Agnihotram. Thyroid dysfunctions in tribal women of Bastar region of Chattisgarh, India. Thyroid Science 2011; 6(6): 1-5.
6. Gharib H., Ryan R.J, Mayberry W.E, and Hockett T., Radioimmunoassay for Triiodothyronine (T3): Affinity and Specificity of Antibody for T3. J Clinical Endocrinol.1971; 33: 509 .
7. Chopra I.J. Ho R. S. and Lam R. An improved Radioimmunoassay of triiodothyronine in human serum. J. Lab Clinical Med. 1971; 80: 729.
8. Sterling L., Diagnosis and Treatment of Thyroid Diseases, Cleveland CRC Press, p. 9-51 (1975).
9. Barker S.B., Determination of Protein Bound Iodine. Journal Biological Chemistry 1948;173:175.
10. Young D. S., Pestaner L.C., and Gilberman U., Effects of Drugs on Clinical Laboratory Tests. Clinical Chemistry 1975; 21: 3660.
11. Nutritive value of Indian foods by C. Gopalan, B.V. Rama Sastri and S.C. Balasubramanian , published by National Institute of Nutrition Indian Council of Medical Research , Hyderabad.
12. Purnima Dey Sarkar, Lincy K. Skaria, Gopinath Agnihotram. Assessment of nutritional status in Hypothyroid patients. J Pharm Biomed Sci. 2013; 27(27):582-585.
13. Arthur JR, Beckett GJ. Thyroid function. British Medical Bulletin 1999; 55: 658-668.
14. Civitareale D, Saiardi A, Falasca P. Purification and characterization of thyroid transcription factor 2. Biochem J 1994;304:981-985.

Conflict of interest: None
Funding: None