

A Study of Thrombocytopenia in Malarial Patients at a Tertiary Care Referral Hospital

Shreya Kavar*, N.S.Jesalpura**, J.M.Shah***

*2nd Year Resident, **Asso. Prof, ***Professor and Head Of Department Of Pathology, AMC MET Medical College & L.G.General Hospital, Ahmedabad.-380008, Gujarat, India.

Abstract: Background & Objective: Malaria is a major public health problem in the tropics with increased morbidity & mortality. It is a protozoal disease caused by parasite of genus plasmodium and transmitted to man by infected female anopheline mosquitos. Anemia and thrombocytopenia are most common. We conducted this study to find out presence and degree of thrombocytopenia. Method: This study was conducted at AMC MET Medical College, L.G. Hospital, Ahmedabad, over a 5 months period. A total 1000 malaria positive cases studied with thick & thin smears stained with Field's stain and platelet count. Results: In the study group of 1000 patients: 700 positive for P. Vivax, 300 positive for P. Falciparum Out of 700 cases of P. Vivax, 600 cases had thrombocytopenia with most cases of Grade 1 & 2 and of 300 cases with P. Falciparum, 250 cases had thrombocytopenia with most cases of Grade 3& 4. So in our study, 85% malaria positive cases had thrombocytopenia. Conclusion: Presence of thrombocytopenia in a patient with acute febrile illness in the tropics increases the probability of malaria. Findings can have therapeutic implication to avoid unnecessary platelet infusion in malarial patients. [Shreya K NJIRM 2017; 8(2):69-71]

Key Words: Malaria, P. Falciparum, P. Vivax, Thrombocytopenia

Author for Correspondence: Shreya R. Kavar, Department Of Pathology, AMC MET Medical College & L. G. Hospital, Ahmedabad-380008. E-Mail: shreyakavar31091@gmail.com M: 9426842818

Introduction: Malaria is a major public health problem in the tropics with increased morbidity and mortality. It is a protozoal disease caused by infection with parasites of the genus Plasmodium and transmitted to man by certain species of infected female anopheline mosquito. Five species of Plasmodium (P. vivax, P.falciparum, P.ovale, P. malariae, P.knowlesi) cause malaria in humans.

From all this five species P.vivax and P.falciparum are the most common. According to World malaria report 2015, there were 214 million cases of malaria with 438000 deaths. The burden was heaviest in WHO African region accounting for 90% of malaria death with more than 2/3rd of death occurring in children aged under 5years¹. Malaria can cause various hematological abnormalities in patients, of which Anemia and Thrombocytopenia are most common. It comprises three distinct stages: Cold, Hot & Sweating. Clinical features vary from mild, moderate & complicated depending on--species of malarial parasite present, patient's state of immunity, Intensity of infection and presence of concomitant conditions. Malarial parasite affects multiple organs of the body such as liver, spleen, brain, gastro intestinal tract, gall bladder, pancreas, blood vessels and placenta. Hence the clinical picture could be of wide spectrum ranging from simple malaise to life threatening central nervous system symptoms like coma^{2,3}.

A number of observational studies have confirmed the association of thrombocytopenia to malaria. Both non-immunological as well as immunological destruction of platelets have been implicated in

causing thrombocytopenia. The speculated mechanisms are coagulation disturbances, sequestration in spleen, antibody mediated platelet destruction, oxidative stress and the role of platelets as cofactors in triggering severe malaria. Abnormalities in platelet structure and function have been described as a consequence of malaria and in rare instances platelets can be invaded by malaria parasites^{4,5,6}.

Aims & Objectives:

We conducted this study to find out the frequency and the degree of thrombocytopenia in patients with malaria, to know the common parasitic species and sex predominance.

Methods: The study was conducted at L.G. Hospital, AMC MET College, Ahmedabad over a five month period. A total 1000 malaria cases positive on peripheral smear were studied. Our study had been approved by ethical committee(IRB).

Malarial parasites were detected by thin & thick smear examination. Thin smear stained by Leishman stain & thick were stained by Field stain. In Field stain polychromatic methylene blue & eosin stains specifically to basophilic & acidophilic cellular elements to demonstrate blood cells and parasites. Absolute platelet count was done by automated analyzer 'XT1800-Sysmex' in hematology section. Data were analyzed and patients with thrombocytopenia have been divided into following five grades:

Grade 0--Within normal limit (1,50,000 or above)

Grade 1--Between 75,000 to 1,50,000

Grade 2-Between 50,000 to 75,000

Grade 3-Between 25,000 to 50,000

Grade 4-Less than 25,000

From this grading, patients were categorized according to parasite infection.

Results: We conducted this study of 1000 malaria positive cases. Result are shown in table 1 and 2.

Most common are grade 1 & grade 2 thrombocytopenia in *P. Vivax* and grade 3 & 4 thrombocytopenia in *P. Falciparum*.

Table-1: Distribution Of Malarial Infection According To Species And Sex

Malarial Infection	No. Of Cases With %	Female (No. Of Cases With %)	Male (No.Of Cases With %)
P.Vivax	700(70%)	288 (41.14%)	412 (58.85%)
P.Falciparum	300 (30%)	102 (34%)	198 (66%)
Total	1000	390 (39%)	610 (61%)

P.vivax is comparatively more common than *P.Falciparum* and it is more common in males.

Table-2: Distribution Of Malarial Infection According To Species And Severity Of Thrombocytopenia

Grading Of Thrombocytopenia	P. Vivax No. Of Cases With %	P. Falciparum No. Of Cases With %
Grade 1 (75,000-1,50,000)	178 (29.66%)	34 (13.6%)
Grade 2 (50,000-75,000)	256 (42.66%)	28 (11.2%)
Grade 3 (25,000-50,000)	110 (18.33%)	128 (51.2%)
Grade 4 (Less Than 25,000)	56 (9.33%)	60 (24%)
Total	600 (100%)	250 (100%)

Discussion: Malaria caused by *P. Vivax* & *P. Falciparum* is endemic in many parts of India. It affects almost all blood components & it is a true hematological disease. Thrombocytopenia & anemia are the most frequent malaria associated hematological complications. In endemic areas malaria has been reported as the major cause of low platelet counts.

15 times^{7,8,9}. Faseela et al.⁵ in her study found similar results. In our study, thrombocytopenia was seen in 85% cases. Colonel et al.¹⁰ reported thrombocytopenia in 72% of patients with malaria infection. Jamal et al.¹¹ in their study on pediatric patient have reported low platelet count in 72% of the patients with malaria infection. However, few studies reported slightly lower incidence of thrombocytopenia like 40%⁶ and 58.97%¹². Exact mechanism of thrombocytopenia in malaria is unknown. Fajardo and Tallent¹³ demonstrated *P. vivax* within platelets and suggested a direct lytic effect of the parasite on the platelets. Both non-immunological destruction as well as immune mechanisms⁷ involving specific platelet associated IgG antibodies that bind directly to malarial antigen in the platelets have been recently reported to play a role in the lysis of platelets¹⁴.

Table -3 Comparison Of Most Common Malarial Species With Other Study

Malarial Species	P. Vivax	P.Falciparum	Total
Present Study	70%	30%	1000
S.J.Khan Study ¹⁸	32%	68%	121
N.K.Gupta Study ¹⁹	56.51%	43.47%	230

This is so characteristic of malaria, that in some places, it is used as an indicator of malaria in patients presenting with fever. Platelets count of less than 150,000/cumm increases the likelihood of malaria 12-

Table-4: Comparison Of Thrombocytopenia Grading Of Our Study With Other Study:-

Malarial Infection	Thrombocytopenia Grade	Present Study	S.J. Khan Study ¹⁸	N.K.Gupta Study ¹⁹
P.Vivax	Grade-1	29.66%	38%	20%
	Grade-2	42.66%	36%	25%
	Grade-3	18.33%	15%	40%
	Grade-4	9.33%	11%	15%
P.Falciparum	Grade-1	13.6%	20%	15.42%
	Grade-2	11.2%	32%	28.57%
	Grade-3	51.2%	36%	50%
	Grade-4	24%	12%	6.0%

Oxidative stress damage of platelets has also been implicated in the etiopathogenesis based on the finding of low levels of platelet superoxide dismutase and glutathione peroxidase activity and high platelet lipid peroxidation levels in malaria patients, when compared to those of healthy subjects¹⁵. Decreased thrombopoiesis has been ruled out, because platelet forming megakaryocytes in the marrow are usually normal or increased^{7,15,16,17}. We found thrombocytopenia in *P.vivax* & *P.falciparum*. Out of which more cases are of *P.vivax* & most severe grade of thrombocytopenia is observed in *P.falciparum*. In S.J.Khan study¹⁸ *P.falciparum* is most common, while in N.K. Gupta study¹⁹ *P.vivax* is most common. In *P.falciparum* Grade 3 Thrombocytopenia Is Common In All Studies.

Conclusion: Presence of thrombocytopenia in a patient with acute febrile illness in tropics increase the probability of malaria. Thrombocytopenia occurs in both malaria but it is more severe in *P. falciparum* as compared to *P.vivax*. Above findings can have therapeutic implication in context of avoiding unnecessary platelet infusion in malarial patients.

References:-

1. WHO / Data and Statistics. www.who.int/malaria/data.
2. Wickramasinghe SN, Abdalla SH. Blood and bone marrow changes in malaria. *Baillieres Best Pract Res Clin Haematol.* 2000;13:277–99.
3. Khan SJ, Khan FR, Usman M, Zahid S. Malaria can lead to thrombocytopenia. *Rawal. Med J.* 2008;33:183–5.
4. Jadhav UM, Patkar VS, Kadam NN. Thrombocytopenia in malaria: Correlation with type and severity of malaria. *J Assoc Physicians India.* 2004;52:615–8.
5. Faseela TS, Roche, Anita KB, Malli CS, Rai Y. Diagnostic value of platelet count in malaria. *J Clin Diagn Res.* 2011;5:464–6.
6. Rasheed A, Saeed S, Khan SA. Platelet counts in malaria. *Pak J Phytopathopathol.* 2008;19:868.
7. Maina RN, Walsh D, Gaddy C, Hongo G, Waitumbi J, Otieno L, et al. Impact of *Plasmodium falciparum* infection on haematological parameters in children living in Western Kenya. *Malar J.* 2010;9(suppl 3):S4.
8. Adedapo AD, Falade CO, Kotila RT, Ademowo GO. Age as a risk factor for thrombocytopenia and anaemia in children treated for acute uncomplicated *falciparum* malaria. *J Vector Borne Dis.* 2007;44:266–71.
9. Lathia TB, Joshi R. Can hematological parameters discriminate malaria from nonmalarious acute febrile illness in the tropics? *Indian J Med Sci.* 2004;58:239–44.
10. Colonel KM, Bhika RD, Khalid S, Khalique-ur-Rehman S, Syes ZA. Severe thrombocytopenia and prolonged bleeding time in patients with malaria (a clinical study of 162 malaria cases) *World Appl Sci J.* 2010;9:484–8.
11. Jamal A, Memon IA, Latif F. The association of *Plasmodium vivax* malaria with thrombocytopenia in febrile children. *Pak Paediatr J.* 2007;31:85–9.
12. Rodríguez-Morales AJ, Sánchez E, Vargas M, Piccolo C, Colina R, Arria M. Anemia and thrombocytopenia in children with *Plasmodium vivax* malaria. *J Trop Pediatr.* 2006;52:49–51.
13. Fajardo LF, Tallent C. Malarial parasites within human platelets. *J Am Med Assoc.* 1974;229:1205–9.
14. Makkar RP, Mukhopadhyay S, Monga A, Gupta AK. *Plasmodium vivax* malaria presenting with severe thrombocytopenia. *Braz J Infect Dis.* 2002;6:263–5.
15. Metanat M, Sharifi-Mood B. Malaria *vivax* and severe thrombocytopenia in Iran. *Iran J Parasitol.* 2010;5:69–70.
16. Gursharan SN, Neha S. Thrombocytopenia and other complications of *Plasmodium vivax* malaria. *Curr Pediatr Res.* 2011;15:117–9.
17. Leowattana W, Tangpukdee N, Thar SK, Nakasiri S, Srivilairit S, Kano S, et al. Changes in platelet count in uncomplicated and severe *falciparum* malaria. *Southeast Asian J Trop Med Public Health.* 2010;41:1035–41.
18. Shiraz Jamal Khan, Yasir Abbas, and Mumtaz Ali Marwat. A Clinical study of thrombocytopenia as an indicator of malaria in adult population. 2012, Article ID 405981
19. Narendra Kumar Gupta, Shyam Babu Bansal, 1 Uttam Chand Jain, 2 and Kiran Sahare 3 *Trop Parasitol.* 2013 Jan-Jun; 3(1): 58–61.

Conflict of interest: None

Funding: None

Cite this Article as: Shreya K, N. Jesalpura, J. Shah. A Study Of Thrombocytopenia In Malarial Patients. *Natl J Integr Res Med* 2017; 8(2):69-71